



## Cytomorphological Study of Breast Lesions in Correlation with Quadrant Location to Assess Prognosis

Dr. Sri Anjani M<sup>1\*</sup>, Dr. Suma Kaza<sup>2</sup>, Dr. Syeda Khadija Fathima<sup>2</sup>, Dr. M Bhavani<sup>3</sup>

<sup>1</sup>Post Graduate, <sup>2</sup>Assistant Professor, <sup>3</sup>Professor & HOD, Department of Pathology, Kamineni Institute of Medical Sciences (KIMS), Akkenepally vari lingotam, Narketpalle, Telangana 508254, India

**ABSTRACT: Introduction:** Tumour location within the breast varies with the highest frequency in the upper outer quadrant (UOQ) and lowest frequency in the lower inner quadrant (LIQ). Whether tumour location is prognostic is unclear. To determine whether tumour location is prognostic, associations between tumour site and clinicopathological characteristics were evaluated. Materials and Methods: All patients enrolled in the Clinical Breast Care Project whose tumour site—UOQ, upper inner quadrant (UIQ), central, LIQ, lower outer quadrant (LOQ)—was determined by a single, dedicated breast pathologist were included in this study. Patients with multicentric disease (n = 73) or tumours spanning multiple quadrants were excluded from further analysis. Clinicopathological characteristics were analysed using chi-square tests for univariate analysis with multivariate analysis performed using principal components analysis (PCA) and multiple logistic regression. Significance was defined as P < 0.05. **Results:** Of the patients with defined tumour location, 30 had bilateral disease. Tumour location in the UOQ (51.5%) was significantly higher than in the UIQ (15.6%), LOQ (14.2%), central (10.6%), or LIQ (8.1%). Tumours in the central quadrant were significantly more likely to have higher tumour stage (P = 0.003) and size (P < 0.001), metastatic lymph nodes (P < 0.001), and mortality (P = 0.011). After multivariate analysis, only tumour size and lymph node status remained significantly associated with survival. **Conclusions:** Evaluation of tumour location as a prognostic factor revealed that although tumours in the central region are associated with less favourable outcome, these associations are not independent of location but rather driven by larger tumour size. Tumours in the central region are more difficult to detect mammographically, resulting in larger tumour size at diagnosis and thus less favourable prognosis. Together, these data demonstrate that tumour location is not an independent prognostic factor.

**Keywords:** Breast Quadrants, Tumour Location, Prognosis.

### RESEARCH PAPER

**\*Corresponding Author:**

Dr. Sri Anjani M  
Post Graduate, Department of Pathology, Kamineni Institute of Medical Sciences (KIMS), Akkenepally vari lingotam, Narketpalle, Telangana 508254, India.

**How to cite this paper:**

Sri Anjani M *et al*;  
“Cytomorphological Study of Breast Lesions in Correlation with Quadrant Location to Assess Prognosis” Middle East Res J. Microbiol Biotechnol., 2021 Nov-Dec 1(1): 6-11.

**Article History:**

| Submit: 27.10.2021 |  
| Revised: 16.11.2021 |  
| Accepted: 29.11.2021 |  
| Published: 27.12.2021 |

**Copyright © 2021 The Author(s):** This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International License (CC BY-NC 4.0) which permits unrestricted use, distribution, and reproduction in any medium for non-commercial use provided the original author and source are credited.

## INTRODUCTION

Importance of quadrant location on prognosis of breast lesions has been investigated for many years. Quadrant location for malignancy has been given importance in (Surveillance Epidemiology End Results) SEER Coding guidelines. However prognostic significance of tumor location in breast cancer remains unclear.

Lymphatic drainage is different for each breast quadrant, therefore absence of axillary node positivity could misclassify high-risk lesion to low risk., Calculated risk for malignant transformation of benign lesions ranges from 3% -17% [1, 2].

Fine needle aspiration cytology (FNAC) is widely accepted as a reliable technique in the initial evaluation of palpable and non-palpable (guided

biopsy) breast lumps. The procedure is simple, safe, cost effective, minimally invasive, rapid, and as sensitive as biopsy [3].

The primary goal of FNAC is to separate malignant lesions that require more radical therapy from benign ones that may be conservatively managed. The scope of cytology now extends into identifying the subtypes of malignant lesions, benign lesions, and minimal residual disease for the purpose of planning the therapeutic protocol and eventual follow-up. Thus, it plays a major role as an important preoperative assessment along with clinical and mammography examination, which together are frequently referred to as “Triple test” [4, 5].

Pattern is identification of quality as a whole. It is known that despite the many sites and many types

of tumors that are aspirated, there are a limited number of patterns based on morphology observed in aspirated material. However, the frequency, significance, and difference of each pattern vary with the site.

Many authors have used various methods to come to a conclusive method of diagnosis on breast lesions. Here, we propose a partially modified method based on systematic pattern analysis to analyze the breast lesions and divide them into individual categories.

Our study focuses on this range of risk by additional consideration of breast quadrant utilizing fine needle aspiration cytology (FNAC) as a diagnostic.

### AIMS AND OBJECTIVES

- To assess the prevalence of breast lesions in various quadrant.
- To assess quadrant of location as a risk factor for breast lesions.
- To correlate cytomorphological features of FNAC of spectrum of lesions encountered with importance of quadrant of location determines the prognosis of the breast lump.

### MATERIALS AND METHODS



One year prospective study done at Department of Pathology in K I M S, Narketpally between June 2017 to July 2018 with 73 cases of breast lump.










Clinical history, radiological imaging and physical examination was done noting quadrant of location along with bilateral examination of nipple, axilla & lymphnodes.

FNAC was done with 22 guage needle. Cytomorphology evaluation carried out on Haematoxylin and Eosin and Giemsa stained smears. Statistics was compiled and p value calculated using SPSS software

Fine needle aspiration was done with a 21- or 22-gauge needle attached to a 10 cc disposable syringe mounted on a syringe holder for single handgrip. The specimen was taken with minimum passes (to minimize hemorrhage) without needle withdrawal and under constant negative

### RESULTS





According to studies conducted by Seth Rummel *et al.*, [10] tumors occurring in lateral upper quadrant carries better prognosis than those occurring in other quadrants. Calculated risk for fibroadenoma is 3%.

*Quadrant wise location of IDCC in both Breast.* The most common location of IDCC in our study was lateral upper quadrant.


RISK FACTORS		LUQ	MUQ
Age at diagnosis	<40 yrs	1	1
	40-49 yrs	4	1
	50 yrs	6	1
Menopausal status	Pre-Menopausal	3	1
	Post-Menopausal	8	2
	Breast feed	9	2
	No	2	1
Tumor size	<2 Cm	0	0
	2-5 Cm	10	3
LN Status	Positive	1	0
	Negative	10	3



The most common location of IDCC in our study was lateral upper quadrant.

## DISCUSSION

Determination of whether tumour location can be used prognostically is important in optimising treatment. Tumour location is highest in the UOQ (50–58%) across multiple populations, including Chinese, Danish, the United Kingdom and women treated within the United States Department of Defence healthcare system. Two studies suggest that the frequency of tumours in the UOQ has increased over time [6, 7]; given the association of tumour in the UOQ with improved prognosis, these data would suggest a trend towards a reduction in breast cancer mortality. Tumour occurrence in the LIQ, however, has also increased significantly and tumour location in the LIQ has been associated with >2-fold increase in mortality, suggesting that the decrease in mortality associated with increasing tumour location in the UOQ may be offset by concomitant increases in the LIQ. In our study, tumours in the UOQ showed a trend towards favourable prognosis although this did not reach the level of

significance ( $P = 0.0754$ ). The less favourable prognosis seen in patients with tumours in the central region can be attributed to increased tumour size and positive lymph node status.<sup>8</sup> Tumour size is thought to reflect the chronological age of the tumour, with smaller tumours being resected earlier than larger tumours. Tumour size has been associated with positive lymph node status in multiple studies. In addition, tumour size is also prognostic in patients with both negative and positive lymph node status. Thus, tumour location in the central region is a surrogate for larger tumour size, resulting in increased rates of metastatic lymph node and breast cancer mortality [21, 22]. Previous studies have demonstrated that tumours within the central region are harder to detect than at other sites and that tumours in this region are more easily detected by clinical examination than mammography. This difficulty in detecting tumours in the central region have been attributed to overpenetration of X-rays in the nipple-areolar complex; accurate diagnosis in this region may require the use of multiple imaging modalities [9, 10, 20].





Pattern recognition is defined as “the act of taking in raw data and taking an action based on category of pattern”. Its aim is to classify patterns based on prior knowledge or statistical information extracted from patterns. Here, the perspective is on identification of pattern that is based on various morphological attributes on aspirates of breast lesions (cellular features, background material, and other features) [11, 12, 19].

Biphasic pattern usually include epithelial cells and myoepithelial cells arranged in mono-layered sheets with a honeycomb pattern (“antler-like, staghorn”). The presence of myoepithelial cells has been recognized as a prominent feature of benign breast disease. Lesions with biphasic pattern are most often benign, hence identification of this pattern is an important aspect of breast lesions [13, 14, 18].

Macrophage-rich pattern is seen predominantly in fibrocystic change and in cysts which usually showed foam cells, apocrine cells, and occasionally non-apocrine cells. Smears with apocrine cells showing degenerative atypia should be interpreted with caution, taking into consideration background patterns like proteinaceous background (for benign cystic lesion of breast) and hemorrhagic background (to rule out malignancy) [15, 16, 17, 23].

## CONCLUSION

The most common age group of breast lesions was 6<sup>th</sup> decade in males and 3<sup>rd</sup> decade in females. There was left breast predominance and lump was the most frequent clinical symptom in both the genders.

Among the benign lesion fibroadenoma (53.4%) showed highest occurrence where as in malignant lesions, it was IDCC (71.4%)

Lateral outer quadrant showed highest involvement for fibroadenoma (58.6%) among the benign lesions Calculated risk for fibroadenoma is very low (3%).

Therefore, it can be concluded that benign lesions occurring in lateral upper quadrant carry good prognosis.

Most common quadrant of involvement for IDCC (78.5) in our study was lateral outer quadrant again. SEER coding for IDCC was 2

Studies have shown IDCC occurring in lateral outer quadrant carries better prognosis when compared to other quadrants. Thus we conclude that, quadrant location of breast lump carries significance in prognosis.

## REFERENCES

1. Nguansangiam, S., Jesdapatarakul, S., & Tangjitgamol, S. (2009). Accuracy of fine needle aspiration cytology from breast masses in Thailand. *Asian Pac J Cancer Prev*, 10(4), 623-6.
2. Kalhan, S., Dubey, S., Sharma, S., & Dudani, S. (2010). Significance of nuclear morphometry in cytological aspirates of breast masses. *Journal of Cytology/Indian Academy of Cytologists*, 27(1), 16-21.
3. Chaiwun, B., Settakorn, J., Ya-in, C., Wisedmongkol, W., Rangdaeng, S., & Thorner, P. (2002). Effectiveness of fine-needle aspiration

- cytology of breast: Analysis of 2,375 cases from northern Thailand. *Diagnostic cytopathology*, 26(3), 201-205.
4. Bhargava, V., Jain, M., Agarwal, K., Thomas, S., & Singh, S. (2008). Critical appraisal of cytological nuclear grading in carcinoma of the breast and its correlation with ER/PR expression. *Journal of Cytology*, 25(2), 58-61.
  5. Joshi, A., & Maimoon, S. (2007). Limitations of fine needle aspiration cytology in subtyping breast malignancies-a report of three cases. *Journal of Cytology*, 24(4), 203-206.
  6. Kaufman, Z., Shpitz, B., Shapiro, M., Rona, R., Lew, S., & Dinbar, A. (1994). Triple approach in the diagnosis of dominant breast masses: combined physical examination, mammography, and fine-needle aspiration. *Journal of surgical oncology*, 56(4), 254-257.
  7. Miller, T. Breast. In: Renshaw, A. A. editor. (2005). *Aspiration cytology – A pattern recognition approach*. 1st ed. China: Elsevier Saunders; 431-76.
  8. Pattari, S. K., Dey, P., Gupta, S. K., & Joshi, K. (2008). Myoepithelial Cells: Any role in aspiration cytology smears of breast tumors?. *CytoJournal*, 5, 9.
  9. Saad, R. S., Silverman, J. F. (2009). Breast. In: Bibbo, M, editor. *Comprehensive cytopathology*. 3rd ed. China: Elsevier Saunders; 713-72.
  10. Rummel, S., Hueman, M. T., Costantino, N., Shriver, C. D., & Ellsworth, R. E. (2015). Tumour location within the breast: Does tumour site have prognostic ability?. *ecancermedicalscience*, 9.
  11. Sohn, V. Y., Arthurs, Z. M., Sebesta, J. A., & Brown, T. A. (2008). Primary tumor location impacts breast cancer survival. *The American journal of surgery*, 195(5), 641-644.
  12. Turner-Warwick, R. T. (1959). The lymphatics of the breast. *British Journal of Surgery*, 46(200), 574-582.
  13. Vendrell-Torne, E., Setoain-Quinquer, J., & Domenech-Torne, F. M. (1972). Study of normal mammary lymphatic drainage using radioactive isotopes. *Journal of Nuclear Medicine*, 13(11), 801-805.
  14. Kurian, A. W., Fish, K., Shema, S. J., & Clarke, C. A. (2010). Lifetime risks of specific breast cancer subtypes among women in four racial/ethnic groups. *Breast Cancer Research*, 12(6), R99.
  15. Farkhanda, J. D., Muhammad, S. A., Ahsan, A. L., Noor, M. K., Imtiaz, S., & Zulfiqar, I. M. (2010). An early diagnosis of benign breast diseases. *J Surg Pak*, 15(4), 74-8.
  16. Godwins, E., David, D., & Akeem, J. (2011). Histopathologic analysis of benign breast diseases in Makurdi, North Central Nigeria. *International Journal of Medicine and Medical Sciences*, 3(5), 125-128.
  17. Haque, T., Khan, G., & Gahlauts, Y. V. S. (1980). Breast lesions: a clinico histopathological study of 200 cases of breast lump. *JAMA*, 150, 1810-1814.
  18. Vijayalakshmi, M., Rao, J. Y., & Shekar, T. Y. (2016). Prevalence of Benign Breast Disease and Risk of Malignancy in Benign Breast Diseases. *IOSR Journal of Dental and Medical Sciences*, 15(8):32-36
  19. Selvakumaran, S., & Sangma, M. B. (2016). Study of various benign breast diseases. *International Surgery Journal*, 4(1), 339-343.
  20. Muddegowda, P. H., Lingegowda, J. B., Kurpad, R., Konapur, P. G., Shivarudrappa, A. S., & Subramaniam, P. M. (2011). The value of systematic pattern analysis in FNAC of breast lesions: 225 cases with cytohistological correlation. *Journal of Cytology/Indian Academy of Cytologists*, 28(1), 13-19.
  21. Chandanwale, S., Rajpal, M., Jadhav, P., Sood, S., Gupta, K., & Gupta, N. (2013). Pattern of benign breast lesions on FNAC in consecutive 100 cases: a study at tertiary care hospital in India. *Int J Pharm Biol Sci*, 3(4), 129-138.
  22. Kamra, H. T., Rana, P., Kaur, S., Verma, S., Munde, S., Ahuja, K., & Singh, K. (2017). Spectrum of Breast Lesions Diagnosed on Fine Needle Aspiration Cytology in Rural Population of Khanpur Kalan, Sonapat (Haryana). *Annals of International Medical and Dental Research*, 3(3), 6-9.
  23. Kroman, N., Wohlfahrt, J., Mouridsen, H. T., & Melbye, M. (2003). Influence of tumor location on breast cancer prognosis. *International journal of cancer*, 105(4), 542-545.
  24. Wu, S., Zhou, J., Ren, Y., Sun, J., Li, F., Lin, Q., ... & He, Z. (2014). Tumor location is a prognostic factor for survival of Chinese women with T1-2N0M0 breast cancer. *International Journal of Surgery*, 12(5), 394-398.